

**2010 Shorebird Banding in Northwest Ohio
Winous Point Shooting Club
Ottawa National Wildlife Refuge
McClure Marsh
H. Thomas Bartlett
Thomas Kashmer**



Tom and Tom holding Ohio's 1st banded Sanderlings

**H. Thomas Bartlett
1833 South Winfield Drive
Tiffin, Ohio 44883
419-447-0005
hthomas.bartlett@gmail.com
Federal Permit #21624**

**Thomas Kashmer
949 County Road 42
Helena, Ohio 43435
419-638-1027
green creekws@gmail.com
Federal Permit #23088**

2010 Shorebird Research Program Report

H. Thomas Bartlett and Thomas Kashmer

Historical

In the summer of 2006, Tom Bartlett and Tom Kashmer began a shorebird banding research project at Winous Point Shooting Club on Sandusky Bay. The purpose of the study was threefold. The first was to determine what species were using northwest Ohio during fall migration and in what numbers. The second was to determine how long they were staying in the area. And the third was, hopefully, to add samples to an avian flu study from shorebirds. During the fall migration of 2006, 226 shorebirds of 10 species plus an additional 10 individuals which were non-shorebirds were banded. Banding occurred on a total of 8 days, with 213.5 net hours of effort, and a respectable 1.11 birds per net hour recorded. During the fall migration of 2007, the effort was repeated at Winous Point. For the 2007 season 281 shorebirds of 12 species plus an additional 23 individuals which were non-shorebirds were banded. Banding occurred on 11 days, with 379.5 net hours of effort, and 0.80 birds per net hour recorded. At the beginning of the 2008 fall migration banding began again at Winous Point. However, early into the season, Doug Brewer, the refuge manager at Ottawa NWR, requested Ottawa NWR be added to the study and with the attempt to collect some avian flu data there as well. The refuge would supply an intern, John Siekierski, to assist our efforts and collect avian flu samples. The research was expanded. For the fall of 2008, there were 40 banding days, 1404.5 net hours of effort, and 1.21 birds per net-hour. 1593 shorebirds of 16 species plus 106 non-shorebirds of 14 species were banded. All total, 1699 individual birds of 30 species were banded during the 2008 season. Several hundred avian flu samples were collected during the process. The fall migration season of 2009 began the same as 2008 with Ottawa NWR and Winous Point Shooting Club as the key banding sites and with the assistance of intern, John Siekierski. Towards the end of the season, a private wetland owned by William and Lena McClure in Sandusky County was added to the study. A total of 1923 individual birds of 46 species were banded at the three sites. These included 1661 individuals of 20 species of shorebirds. Banding occurred over 51 days and included 1808 net hours of effort for 1.06 birds banded per net hour. Over the four seasons, 21 species of shorebirds of 3761 individuals have been banded, plus an additional 400 non-shorebirds. A total 110 days with 3805.5 net hours of effort have resulted in 0.99 shorebirds banded per net hour. In addition, numerous birds were recaptured from previous seasons to demonstrate the importance of the western Lake Erie wetlands to migrating shorebirds. Again, several hundred avian flu samples were collected and given to Dr. Richard Slemmons of Ohio State University. All samples were negative for the dangerous strains of avian flu but several less dangerous strains were found.

2010 Summary

Beginning July 15, 2010 the shorebird research program continued into its fifth season. Again, support of money, personnel, and facilities were supplied by Winous Point Shooting Club/Winous Point Conservancy and Ottawa National Wildlife Refuge. Winous Point supplied banding sites, funding for new equipment and personnel for setup and moving nets. Ottawa National Wildlife Refuge supplied banding sites, funding for nets, equipment, and personnel. In addition to intern, John Siekierski, several other summer interns were supplied at peak times to assist in the capture, sampling, and banding of shorebirds. Again this year Mr. Siekierski collected hundreds of samples from shorebirds for the avian flu study database for Dr. Richard Slemmons at Ohio State University. And for the fifth year of sampling shorebirds, no positive results of avian flu was found. In addition, the new 2009 site, the property of William and Lena McClure, in Rice Township, Section 30 of Sandusky County about a mile south of the Winous Point properties was also used. Again recapture data shows that the shorebirds are interacting between all three sites.

The 2010 banding project continued to improve as it has for each of the years of the study. This year banding occurred on 46 days with several of those days at two sites at the same time. Net hours of banding effort also increased to 1808. This brings the total banding days to 156 days over five years and 5513.5 net hours of effort. This year 1529 shorebirds of 20 species were banded. Of the twenty species of shorebirds, seven species are of note. Ohio's first Sanderlings(2), first Baird's Sandpipers(4) and first Buff-breasted Sandpipers(3) were banded. Four other species which are rarely banded in Ohio were Marbled Godwit, Western Sandpiper, White-rumped Sandpiper, and 15 Red-necked Phalarope. In addition, 222 non-shorebirds of 26 species were banded. Unusual among these were Hooded Merganser, Sora, and Northern Waterthrush. The total number of birds banded in 2010 was 1751. 200 individual recaptures of 177 individuals, 6 escape/released individuals, and 8 casualties were recorded. Of the recaptured birds, 1 was banded in 2007, 7 were banded in 2008, and 7 were banded in 2009. The rest were all banded early in the season. The average length of time between banding and recapture was 5.8 days with a range of 1 to 26 days and an average weight gain of 13.38% body mass. Of the recaptured individuals, 70% showed a weight gain at recapture, 27% showed a weight loss at recapture, and 3% show no change in weight at recapture. A more statistical analysis of the recapture data will be done. There appears to be a relationship between the length of time after capture and weight gain. The actual process of capture and banding may have a short-term negative effect. The casualty rate was a very good 0.46% of birds banded and 0.41% of birds handled. All casualties were deposited at the Cleveland Museum of Natural History.

A breakdown of all the data is in the following pages. Electronic copies of all data collected will be sent to all supporters of this research. With the inclusion of the 2010 data, we have now banded 5822 individuals. Of these 5200 were shorebirds of 24 species. Project goals are clearly being met.

Shorebird Banding Results Shorebirds vs. Non-shorebirds Banded

2006

Shorebirds	226 (96%)
Non-shorebirds	10 (4%)
Total	236

2007

Shorebirds	281 (93%)
Non-shorebirds	22 (7%)
Total	303

Two-year Totals

(507 – 94%)
(32 – 6%)
(539)

2008

Shorebirds	1593 (94%)
Non-shorebirds	106 (6%)
Total	1699

Three-year Totals

(2100 – 94%)
(138 – 6%)
(2238)

2009

Shorebirds	1661 (86%)
Non-shorebirds	262 (14%)
Total	1923

Four-year Totals

(3671 – 88%)
(400 – 12%)
(4161)

2010

Shorebirds	1529(87%)
Non-shorebirds	222 (13%)
Total	1751

Five-year Totals

(5200 – 89%)
(622 – 11%)
(5822)

2010 Totals per Banding Site

Species	Location	ONWR	Winous	McClure	Total	
Semipalmated Plover		4	30	5	39	
Killdeer		62	8	48	118	
Spotted Sandpiper		5		4	9	
Solitary Sandpiper		3		3	6	
Lesser Yellowlegs		13	43	51	107	
Marbled Godwit			1		1	
Sanderling		2			2	
Semipalmated Sandpiper		18	300	157	475	
Western Sandpiper			2		2	
Least Sandpiper		179	124	151	454	
White-rumped Sandpiper			1		1	
Baird's Sandpiper			4		4	
Pectoral Sandpiper		10	28	96	134	
Dunlin		6			6	
Stilt Sandpiper		3	8	3	14	
Buff-breasted Sandpiper			3		3	
Short-billed Dowitcher		4	105	25	134	
Long-billed Dowitcher			1		1	
American Woodcock			*		*	
Wilson's Snipe		3	1		4	
Red-necked Phalarope		1	14		15	
Total species -shorebirds	20	14	16	10	20	
Total individuals - shorebirds		313	673	543	1529	
Wood Duck				1	1	
Hooded Merganser				1	1	
Sora		1			1	
"Trail's" Flycatcher		4			4	
Eastern Kingbird		2			2	
Purple Martin			1	4	5	
Tree Swallow		2	16	10	28	
Northern Rough-winged Swallow				7	7	
Bank Swallow			18	7	25	
Barn Swallow		5	8	16	29	
Marsh Wren		2			2	
American Robin		1			1	
European Starling		4		2	6	
Yellow Warbler		4			4	
Northern Waterthrush		1	1		2	
Common Yellowthroat				1	1	
Savannah Sparrow		1	1		2	
Song Sparrow		9	3	1	13	
Indigo Bunting		1	1	1	3	
Red-winged Blackbird		16	16	33	65	
Common Grackle		1		1	2	
Baltimore Oriole		1			1	
American Goldfinch		1	1	15	17	
Total species - nonshorebirds	23	17	10	14	23	
Total individuals - nonshorebirds		56	66	100	222	
Total birds		369	739	643	1751	
Total species banded		31	26	24	43	
Banding Days	46	13	21	14	48	(2 days at 2 sites)
Net hours		377.25	624.75	706	1708	
Bird per net hour		0.98	1.18	0.91	1.03	
Recaptures		22	144	34	200	(177 individuals)
Escapes/Releases		5	1	0	6	
Casulties		0	5	3	8	

* = An American Woodcock was captured and banded near the WPSC offices on 8/5/2010
This was during the Wild Thing Day banding demonstration



Bartlett with Marbled Godwit at Winous Point Point



Buff-breasted Sandpiper at Winous



Dueling Baird's Sandpipers



White-rumped Sandpiper